



Plant Risk Evaluator -- PRE Evaluation Report

Buddleja davidii 'Black Knight' -- Illinois

2017 Farm Bill PRE Project

PRE Score: 16 -- Reject (high risk of invasiveness)

Confidence: 70 / 100

Questions answered: 20 of 20 -- Valid (80% or more questions answered)

Privacy: Public Status: Submitted

Evaluation Date: April 30, 2017

This PDF was created on June 15, 2018

Plant Evaluated

Buddleja davidii 'Black Knight'



Image by Missouri Botanical Garden

Evaluation Overview

A PRETM screener conducted a literature review for this plant ($Buddleja\ davidii\ 'Black\ Knight'$) in an effort to understand the invasive history, reproductive strategies, and the impact, if any, on the region's native plants and animals. This research reflects the data available at the time this evaluation was conducted.

General Information

Status: Submitted

Screener: Emily Russell

Evaluation Date: April 30, 2017

Plant Information

Plant: Buddleja davidii 'Black Knight'

If the plant is a cultivar, how does its behavior differs from its parent's?

Black Knight has darker flowers than the species.

Regional Information

Region Name: Illinois

Climate Matching Map

To answer four of the PRE questions for a regional evaluation, a climate map with three climate data layers (Precipitation, UN EcoZones, and Plant Hardiness) is needed. These maps were built using a toolkit created in collaboration with GreenInfo Network, USDA, PlantRight, California-Invasive Plant Council, and The Information Center for the Environment at UC Davis.

Click <u>here</u> to see the generated climate matching map for this region. This climate match database is hosted by GreenInfo Network and publicly accessible.

Evaluation Questions

These questions are based in an original article published at the University of California, Davis, and can be found on the PLOS One website, here: https://doi.org/10.1371/journal.pone.0121053

Invasive History and Climate Matching (Questions 1 - 6)

- 1. Has the species (or cultivar or variety, if applicable; applies to subsequent "species" questions) become naturalized where it is not native?
 - Answer: Yes, which contributes 1 points to the total PRE score.
 - The *screener* has a **Medium** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja davidii has naturalized in North and South America, Europe, Africa, and Australia. There is no evidence that Buddleja davidii 'Black Knight' is significantly different than the species in reproductive traits and ability to naturalize. "B. davidii, by nature, establishes on frequently disturbed sites which provide source for satellite populations that can repopulate new and post-disturbance landscapes. As it is well known that species invasion via multiple loci is the most effective means of establishing non-indigenous species in a new environment (Mack et al., 2000), the continued presence of non-sterile B. davidii cultivars in urban, residential, and wild areas guarantees continued reintroduction into native areas (regardless of native control efforts)." (Tallent-Halsell)

Reference(s):

- Randall, R. (2012). A Global Compendium of Weeds. 2nd Edition..
- Ebeling, S., & Tallent-Halsell N. (2009). Buddleja davidii (butterfly bush) Datasheet In: Invasive Species Compendium.
- USDA NRCS (2016). USDA PLANTS Database: Buddleja davidii (orange eye butterflybush).

2. Is the species (or cultivar or variety) noted as being naturalized in the US or world in a similar climate?

- Answer: Yes, which contributes 2 points to the total PRE score.
- The *screener* has a **Medium** confidence in this answer based on the available literature.

Buddleja davidii has naturalized in Korea, the eastern United States, and areas of eastern and northern Europe where there is a similar climate to Illinois. There is no evidence that Buddleja davidii 'Black Knight' is significantly different than the species in reproductive traits and ability to naturalize.

Reference(s):

- Ebeling, S., & Tallent-Halsell N. (2009). Buddleja davidii (butterfly bush) Datasheet In: Invasive Species Compendium.
- USDA NRCS (2016). USDA PLANTS Database: Buddleja davidii (orange eye butterflybush).

3. Is the species (or cultivar or variety) noted as being invasive in the U.S. or world?

- Answer: Yes, which contributes 2 points to the total PRE score.
- The *screener* has a **Medium** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja davidii is classified as a quarantine weed in Oregon (except ODA approved sterile cultivars) and a noxious weed in Washington. In New Zealand it is estimated that the weed pest is displacing valued native species and costing the forestry industry \$0.5 to 2.9 million annually in control and lost production (USFS). It is also invasive in the United Kingdom. There is no evidence that Buddleja davidii 'Black Knight' is significantly different than the species in reproductive traits and ability to invade natural habitats.

- Squirrell, J. (2015). Buddleia, Buddleja davidii GB non-native species secretariat.
- USDA Forest Service Forest Health Staff (2005). Weed of the Week: Butterfly Bush (Buddleja davidii Franch.).
- Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.
- Ebeling, S., & Tallent-Halsell N. (2009). Buddleja davidii (butterfly bush) Datasheet In: Invasive Species Compendium.
- USDA NRCS (2016). USDA PLANTS Database: Buddleja davidii (orange eye butterflybush).

4. Is the species (or cultivar or variety) noted as being invasive in the US or world in a similar climate?

- Answer: No, which contributes 0 points to the total PRE score.
- The *screener* has a **Low** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja davidii has not been documented as an invasive species in a climate similar to Illinois. It is on a few invasive lists in the Mid-Atlantic as a species to watch.

Reference(s):

• Swearingen, J., Slattery B., Reshetiloff K., & Zwicker S. (2010). Plant Invaders of Mid-Atlantic Natural Areas. 168.

5. Are other species of the same genus (or closely related genera) invasive in a similar climate?

- Answer: No, which contributes 0 points to the total PRE score.
- The *screener* has a **Low** confidence in this answer based on the available literature.

Answer / Justification:

There are 23 other species of Buddleja in the Global Compendium of weeds although B. davidii has by far the most number of citations. Buddleja generally seems to be invasive in climates that are warmer than Illinois.

Reference(s):

• Randall, R. (2012). A Global Compendium of Weeds. 2nd Edition..

6. Is the species (or cultivar or variety) found predominately in a climate matching the region of concern?

- Answer: No, which contributes 0 points to the total PRE score.
- The *screener* has a **High** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja davidii 'Black Knight' is widely cultivated and will grow in many different climates.

Reference(s):

• GBIF Secretariat (2016). GBIF Backbone Taxonomy: Buddleja davidii Franch...

Impact on Native Plants and Animals (Questions 7 - 10)

- 7. Does this plant displace native plants and dominate (overtop or smother) the plant community in areas where it has established?
 - Answer: Yes, which contributes 1 points to the total PRE score.
 - The *screener* has a **High** confidence in this answer based on the available literature.

Answer / Justification:

"Buddleja davidii out-competes native floras producing dense monotypic thickets." The cultivar 'Black Knight' is not significantly different than the species in habit or reproduction.

- Squirrell, J. (2015). Buddleia, Buddleja davidii GB non-native species secretariat.
- Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.

8. Is the plant noted as promoting fire and/or changing fire regimes?

- Answer: No, which contributes 0 points to the total PRE score.
- The *screener* has a **Low** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja davidii is not noted as promoting fire or changing fire regimes.

Reference(s):

• [Anonymous].

9. Is the plant a health risk to humans or animals/fish? Has the species been noted as impacting grazing systems?

- Answer: No, which contributes 0 points to the total PRE score.
- The *screener* has a **Medium** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja davidii is palatable to cattle and goats and has evolved strategies to survive defoliation, but there is no evidence that it impacts grazing systems.

Reference(s):

• Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.

10. Does the plant produce impenetrable thickets, blocking or slowing movement of animals, livestock, or humans?

- Answer: Yes, which contributes 1 points to the total PRE score.
- The *screener* has a **High** confidence in this answer based on the available literature.

"Buddleja davidii out-competes native floras producing dense monotypic thickets." The cultivar 'Black Knight' is not significantly different than the species in habit or reproduction.

Reference(s):

- Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.
- Squirrell, J. (2015). Buddleia, Buddleja davidii GB non-native species secretariat.

Reproductive Strategies (Questions 11 - 17)

11. Does this species (or cultivar or variety) reproduce and spread vegetatively?

- Answer: Yes, which contributes 1 points to the total PRE score.
- The *screener* has a **High** confidence in this answer based on the available literature.

Answer / Justification:

"Plants readily reproduce asexually from stem and root fragments (Miller, 1984; Smale, 1990). B. davidii individuals that have been disturbed by flooding and mechanical means have been observed regenerating from buried stems, stumps, and roots soon after the disruption (N. Tallent-Halsell, unpubl.). B. davidii debris, left after removal attempts, can regenerate, flower, and spread, if left in on site on floodplains (H. Turnbull, pers. comm.)"

Reference(s):

• Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.

12. If naturally detached fragments from this plant are capable of producing new plants, is this a common method of reproduction for the plant?

- Answer: Yes, which contributes 1 points to the total PRE score.
- The *screener* has a **High** confidence in this answer based on the available literature.

"Plants readily reproduce asexually from stem and root fragments (Miller, 1984; Smale, 1990). B. davidii individuals that have been disturbed by flooding and mechanical means have been observed regenerating from buried stems, stumps, and roots soon after the disruption (N. Tallent-Halsell, unpubl.). B. davidii debris, left after removal attempts, can regenerate, flower, and spread, if left in on site on floodplains (H. Turnbull, pers. comm.)"

Reference(s):

• Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.

13. Does the species (or cultivar or variety) commonly produce viable seed?

- Answer: Yes, which contributes 1 points to the total PRE score.
- The *screener* has a **Very High** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja davidii commonly reproduces by seed. In a Florida study, Buddleja davidii 'Black Knight' had an average of 281,000 to 973,000 seeds per plant.

Reference(s):

• Wilson, S. B., Mecca L. K., Gersony J. A., Thetford M., & Raymer J. S. (2004). Evaluation of 14 Butterfly Bush Taxa Grown in Western and Southern Florida: II. Seed Production and Germination. HortTechnology. 14, 612–618.

14. Does this plant produce copious viable seeds each year (> 1000)?

- Answer: Yes, which contributes 1 points to the total PRE score.
- The screener has a Very High confidence in this answer based on the available literature.

"A single mature Buddleja davidii individual can produce millions of seeds." In a Florida study, Buddleja davidii 'Black Knight' had an average of 281,000 to 973,000 seeds per plant, with a germination rate of 14.7-28.6%.

Reference(s):

- Wilson, S. B., Mecca L. K., Gersony J. A., Thetford M., & Raymer J. S. (2004). Evaluation of 14 Butterfly Bush Taxa Grown in Western and Southern Florida: II. Seed Production and Germination. HortTechnology. 14, 612–618.
- Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.

15. Is there significant germination (>25%) of seeds the next growing season, with no requirement of an infrequent environmental condition for seeds to germinate (i.e. fire) or long dormancy period?

- Answer: Yes, which contributes 1 points to the total PRE score.
- The *screener* has a **Medium** confidence in this answer based on the available literature.

Answer / Justification:

In a Florida study, Buddleja davidii 'Black Knight' had an average of 281,000 to 973,000 seeds per plant, with a germination rate of 14.7-28.6%. According to this study, Buddleja seeds have the capacity to germinate readily over a broad range of temperatures, but will not germinate in the dark.

- Wilson, S. B., Mecca L. K., Gersony J. A., Thetford M., & Raymer J. S. (2004). Evaluation of 14 Butterfly Bush Taxa Grown in Western and Southern Florida: II. Seed Production and Germination. HortTechnology. 14, 612–618.
- Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.

16. Does this plant produce viable seed within the first three years (for an herbaceous species) to five years (for a woody species) after germination?

- Answer: Yes, which contributes 1 points to the total PRE score.
- The *screener* has a **High** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja davidii typically flowers at two years of age.

Reference(s):

• Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.

17. Does this plant continuously produce seed for >3 months each year or does seed production occur more than once a year?

- Answer: Yes, which contributes 1 points to the total PRE score.
- The *screener* has a **Very High** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja davidii 'Black Knight' flowers from May through October in Illinois. Seed formation and ripening typically occurs within 3 weeks after flowering.

- Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.
- Chicago Botanic Garden (2017). Chicago Botanic Garden Plant Finder.

Dispersal (Questions 18 - 20)

18. Are the plant's propagules frequently dispersed long distance (>100 m) by mammals or birds or via domestic animals?

- Answer: No, which contributes 0 points to the total PRE score.
- The *screener* has a **Medium** confidence in this answer based on the available literature.

Answer / Justification:

There are not reports of dispersal by animals.

Reference(s):

• [Anonymous].

19. Are the plant's propagules frequently dispersed long distance (>100 m) by wind or water?

- Answer: **Yes**, which contributes **1** points to the total PRE score.
- The *screener* has a **Medium** confidence in this answer based on the available literature.

Answer / Justification:

Water could disperse Buddleja seeds long distances along sea coasts, flood plains, and riparian corridors. Buddleja davidii 'Black Knight' has very fine seeds inside a winged sheath, likely to aid in dispersal by wind. Maximum wind dispersal distance has not been determined, but is likely to be less than 100m. "Our results show that normal wind dispersal occurs on a scale (3 m-15 m) that is insufficient to explain the establishment of isolated populations. Alternative dispersal methods—unusually high wind events, birds, mammals, boots, and tires—require further study to explain butterfly bush long-distance invasion in non-riparian sites."

Reference(s):

- Wilson, S. B., Mecca L. K., Gersony J. A., Thetford M., & Raymer J. S. (2004). Evaluation of 14 Butterfly Bush Taxa Grown in Western and Southern Florida: II. Seed Production and Germination. HortTechnology. 14, 612–618.
- Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.
- Ream, J. (2006). Production and invasion of Butterfly bush (Buddleja davidii) in Oregon.

20. Are the plant's propagules frequently dispersed via contaminated seed (agriculture or wildflower packets), equipment, vehicles, boats or clothing/shoes?

- Answer: Yes, which contributes 1 points to the total PRE score.
- The *screener* has a **High** confidence in this answer based on the available literature.

Answer / Justification:

Buddleja seeds are dispersed by cars, trains, and equipment.

Reference(s):

• Tallent-Halsell, N. G., & Watt M. S. (2009). The Invasive Buddleja davidii (Butterfly Bush). Botanical Review. 75, 292–325.

Total PRE Score

PRE Score: 16 -- Reject (high risk of invasiveness)

Confidence: 70 / 100

Questions answered: 20 of 20 -- Valid (80% or more questions answered)

PRE Score Legend

The PRE Score is calculated by adding the point totals for each (answered) question.

< 13 : accept (low risk of invasiveness)

13 - 15 : evaluate further

> 15 : reject (high risk of invasiveness)

Questions Answered Legend

It is important to answer at least 16 questions to consider a PRE Score as "valid".

>= 16 : valid (80% or more questions answered)

<= 15 : invalid (not enough questions answered)

Organization Ownership and Content Privacy

Organization: 2017 Farm Bill PRE Project

Content Privacy: Public

Evaluation Reviewers

The PRE approach is to base decisions on science and make decisions by consensus of diverse horticultural stakeholders. The literature review and process of answering PRE's questions are based on science; the decisions of which plants to prioritize are based on consensus. To ensure this process is in place and that PRE is collaborative, volunteer stakeholders are recruited from each region to review evaluations. The following experts in their profession (plant science, conservation, or horticultural trade) have participated as volunteer PRE reviewers for this evaluation:

This evaluation does not have any reviewers.

Evaluation Issues

The following section lists all public issues for this evaluation. Issues provide a way for stakeholder reviewers to communicate any concerns or suggestions they might have with the plant or evaluation. Please email PlantRight@suscon.org if additional action is required to resolve open issues.

There are currently no issues associated with this evaluation.

About PRE and this Plant Evaluation Report

The PlantRight Plant Risk Evaluator -- PRE is an online database and platform enabling those involved in non-native, terrestrial plant production to know before they grow if a plant poses a regional invasive risk. This tool offers many benefits, and we encourage you to visit the PRE website (https://pre.ice.ucdavis.edu) for more information.

If you are a nursery trade association, or involved in the research, development or distribution of horticultural plants we invite you to join the PRE community. If you are a plant scientist, affiliated with a horticultural college or botanic garden, and would like to learn more about becoming a PRE Screener, please drop us an email, PlantRight@suscon.org, requesting a PRE Account.

PRE beta funding is provided by Sustainable Conservation (http://www.suscon.org/) and a USDA Farm Bill grant.